

A NOTE ON INVARIANT DIFFERENTIAL OPERATORS ON SIEGEL-JACOBI SPACE

JAE-HYUN YANG

ABSTRACT. For two positive integers m and n , we let $\mathbb{H}_n$ be the Siegel upper half plane of degree n and let $\mathbb{C}^{(m,n)}$ be the set of all $m \times n$ complex matrices. In this article, we investigate differential operators on the Siegel-Jacobi space $\mathbb{H}_n \times \mathbb{C}^{(m,n)}$ that are invariant under the natural action of the Jacobi group $Sp(n, \mathbb{R}) \ltimes H_{\mathbb{R}}^{(n,m)}$ on $\mathbb{H}_n \times \mathbb{C}^{(m,n)}$, where $H_{\mathbb{R}}^{(n,m)}$ denotes the Heisenberg group.

1. Introduction

For a given fixed positive integer n , we let

$$\mathbb{H}_n = \{ \Omega \in \mathbb{C}^{(n,n)} \mid \Omega = {}^t\Omega, \quad \text{Im } \Omega > 0 \}$$

be the Siegel upper half plane of degree n and let

$$Sp(n, \mathbb{R}) = \{ M \in \mathbb{R}^{(2n,2n)} \mid {}^tM J_n M = J_n \}$$

be the symplectic group of degree n , where $F^{(k,l)}$ denotes the set of all $k \times l$ matrices with entries in a commutative ring F for two positive integers k and l , tM denotes the transpose matrix of a matrix M and

$$J_n = \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix}.$$

$Sp(n, \mathbb{R})$ acts on $\mathbb{H}_n$ transitively by

$$(1.1) \quad M \cdot \Omega = (A\Omega + B)(C\Omega + D)^{-1},$$

where $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in Sp(n, \mathbb{R})$ and $\Omega \in \mathbb{H}_n$.

For two positive integers m and n , we consider the Heisenberg group

$$H_{\mathbb{R}}^{(n,m)} = \{ (\lambda, \mu; \kappa) \mid \lambda, \mu \in \mathbb{R}^{(m,n)}, \kappa \in \mathbb{R}^{(m,m)}, \kappa + \mu {}^t\lambda \text{ symmetric} \}$$

endowed with the following multiplication law

$$(\lambda, \mu; \kappa) \circ (\lambda', \mu'; \kappa') = (\lambda + \lambda', \mu + \mu'; \kappa + \kappa' + \lambda {}^t\mu' - \mu {}^t\lambda').$$

We define the semidirect product of $Sp(n, \mathbb{R})$ and $H_{\mathbb{R}}^{(n,m)}$

$$G^J = Sp(n, \mathbb{R}) \ltimes H_{\mathbb{R}}^{(n,m)}$$

2000 Mathematics Subject Classification. Primary 13A50, 32Wxx, 15A72.

Keywords and phrases: invariants, invariant differential operators, Siegel-Jacobi space.

This work was supported financially by Inha University Grant.

endowed with the following multiplication law

$$(M, (\lambda, \mu; \kappa)) \cdot (M', (\lambda', \mu'; \kappa')) = (MM', (\tilde{\lambda} + \lambda', \tilde{\mu} + \mu'; \kappa + \kappa' + \tilde{\lambda}^t \mu' - \tilde{\mu}^t \lambda'))$$

with $M, M' \in Sp(n, \mathbb{R})$, $(\lambda, \mu; \kappa), (\lambda', \mu'; \kappa') \in H_{\mathbb{R}}^{(n,m)}$ and $(\tilde{\lambda}, \tilde{\mu}) = (\lambda, \mu)M'$. Then G^J acts on $\mathbb{H}_n \times \mathbb{C}^{(m,n)}$ transitively by

$$(1.2) \quad (M, (\lambda, \mu; \kappa)) \cdot (\Omega, Z) = \left(M \cdot \Omega, (Z + \lambda\Omega + \mu)(C\Omega + D)^{-1} \right),$$

where $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in Sp(n, \mathbb{R})$, $(\lambda, \mu; \kappa) \in H_{\mathbb{R}}^{(n,m)}$ and $(\Omega, Z) \in \mathbb{H}_n \times \mathbb{C}^{(m,n)}$.

We note that the Jacobi group G^J is *not* a reductive Lie group and also that the space $\mathbb{H}_n \times \mathbb{C}^{(m,n)}$ is not a symmetric space. We refer to [1, 3, 16-19, 21-25] about automorphic forms on G^J and topics related to the content of this paper. From now on, for brevity we write $\mathbb{H}_{n,m} = \mathbb{H}_n \times \mathbb{C}^{(m,n)}$.

The aim of this paper is to study differential operators on $\mathbb{H}_{n,m}$ which are invariant under the action (1.2) of G^J . This article is organized as follows. In Section 2, we review differential operators on $\mathbb{H}_n$ invariant under the action (1.1) of $Sp(n, \mathbb{R})$. We let $\mathbb{D}(\mathbb{H}_n)$ denote the algebra of all differential operators on $\mathbb{H}_n$ that are invariant under the action (1.1). We briefly describe the work of Maass [10] about constructing explicit algebraically independent generators of $\mathbb{D}(\mathbb{H}_n)$ and Shimura's construction [12] of canonically defined algebraically independent generators of $\mathbb{D}(\mathbb{H}_n)$. In Section 3, we investigate differential operators on $\mathbb{H}_{n,m}$ invariant under the action (1.2) of G^J . For two positive integers m and n , we let

$$T_{n,m} = \{ (\omega, z) \mid \omega = {}^t\omega \in \mathbb{C}^{(n,n)}, z \in \mathbb{C}^{(m,n)} \}$$

be the complex vector space of dimension $\frac{n(n+1)}{2} + mn$. We have a *natural action* of the unitary group $U(n)$ on $T_{n,m}$ given by

$$(1.3) \quad u \cdot (\omega, z) = (u \omega {}^t u, z {}^t u), \quad u \in U(n), (\omega, z) \in T_{n,m}.$$

The action (1.3) of $U(n)$ induces naturally the representation τ of $U(n)$ on the polynomial algebra $\text{Pol}(T_{n,m})$ consisting of complex valued polynomial functions on $T_{n,m}$. We show that the study of differential operators on $\mathbb{H}_{n,m}$ invariant under the action (1.2) of G^J is equivalent to that of polynomials on $T_{n,m}$ that are invariant under the representation τ of $U(n)$ on $\text{Pol}(T_{n,m})$. Let $\text{Pol}(T_{n,m})^{U(n)}$ denote the subalgebra of $\text{Pol}(T_{n,m})$ consisting of all polynomials on $T_{n,m}$ invariant under the representation τ of $U(n)$, and $\mathbb{D}(\mathbb{H}_{n,m})$ denote the algebra of all differential operators on $\mathbb{H}_{n,m}$ invariant under the action (1.2) of G^J . We note that $\mathbb{D}(\mathbb{H}_{n,m})$ is *not* commutative. The main important problem is to find explicit algebraically independent generators of $\text{Pol}(T_{n,m})^{U(n)}$ and algebraically independent generators of $\mathbb{D}(\mathbb{H}_{n,m})$. We provides many $U(n)$ -invariants in $\text{Pol}(T_{n,m})^{U(n)}$ which may contain algebraically independent generators of $\text{Pol}(T_{n,m})^{U(n)}$. We also provides some explicit G^J -invariant differential operators on $\mathbb{H}_{n,m}$ that are obtained by complicated computations. As an example, we give explicit algebraically independent generators of $\mathbb{D}(\mathbb{H}_{1,1})$. We want to mention

that at this moment it is quite complicated and difficult to find the generators of the algebra of all invariant differential operators on $\mathbb{H}_{n,m}$ and to express invariant differential operators on $\mathbb{H}_{n,m}$ explicitly.

2. Invariant differential operators on $\mathbb{H}_n$

For $\Omega = (\omega_{ij}) \in \mathbb{H}_n$, we write $\Omega = X + iY$ with $X = (x_{ij})$, $Y = (y_{ij})$ real. We put $d\Omega = (d\omega_{ij})$ and $d\bar{\Omega} = (d\bar{\omega}_{ij})$. We also put

$$\frac{\partial}{\partial \Omega} = \left(\frac{1 + \delta_{ij}}{2} \frac{\partial}{\partial \omega_{ij}} \right) \quad \text{and} \quad \frac{\partial}{\partial \bar{\Omega}} = \left(\frac{1 + \delta_{ij}}{2} \frac{\partial}{\partial \bar{\omega}_{ij}} \right).$$

Then for a positive real number A ,

$$(2.1) \quad ds_{n,A}^2 = A \operatorname{tr} \left(Y^{-1} d\Omega Y^{-1} d\bar{\Omega} \right)$$

is a $Sp(n, \mathbb{R})$ -invariant Kähler metric on $\mathbb{H}_n$ (cf. [13, 14]), where $\operatorname{tr}(M)$ denotes the trace of a square matrix M . H. Maass [9] proved that the Laplacian of $ds_{n,A}^2$ is given by

$$(2.2) \quad \Delta_{n,A} = \frac{4}{A} \operatorname{tr} \left(Y^t \left(Y \frac{\partial}{\partial \bar{\Omega}} \right) \frac{\partial}{\partial \Omega} \right).$$

And

$$dv_n(\Omega) = (\det Y)^{-(n+1)} \prod_{1 \leq i \leq j \leq n} dx_{ij} \prod_{1 \leq i \leq j \leq n} dy_{ij}$$

is a $Sp(n, \mathbb{R})$ -invariant volume element on $\mathbb{H}_n$ (cf. [14, p. 130]).

For brevity, we write $G = Sp(n, \mathbb{R})$. The isotropy subgroup K at iI_n for the action (1.1) is a maximal compact subgroup given by

$$K = \left\{ \begin{pmatrix} A & -B \\ B & A \end{pmatrix} \mid A^t A + B^t B = I_n, \ A^t B = B^t A, \ A, B \in \mathbb{R}^{(n,n)} \right\}.$$

Let $\mathfrak{k}$ be the Lie algebra of K . Then the Lie algebra $\mathfrak{g}$ of G has a Cartan decomposition $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$, where

$$\mathfrak{p} = \left\{ \begin{pmatrix} X & Y \\ Y & -X \end{pmatrix} \mid X = {}^t X, \ Y = {}^t Y, \ X, Y \in \mathbb{R}^{(n,n)} \right\}.$$

The subspace $\mathfrak{p}$ of $\mathfrak{g}$ may be regarded as the tangent space of $\mathbb{H}_n$ at iI_n . The adjoint representation of G on $\mathfrak{g}$ induces the action of K on $\mathfrak{p}$ given by

$$(2.3) \quad k \cdot Z = kZ {}^t k, \quad k \in K, \ Z \in \mathfrak{p}.$$

Let T_n be the vector space of $n \times n$ symmetric complex matrices. We let $\Psi : \mathfrak{p} \longrightarrow T_n$ be the map defined by

$$(2.4) \quad \Psi \left(\begin{pmatrix} X & Y \\ Y & -X \end{pmatrix} \right) = X + iY, \quad \begin{pmatrix} X & Y \\ Y & -X \end{pmatrix} \in \mathfrak{p}.$$

We let $\delta : K \longrightarrow U(n)$ be the isomorphism defined by

$$(2.5) \quad \delta \left(\begin{pmatrix} A & -B \\ B & A \end{pmatrix} \right) = A + iB, \quad \begin{pmatrix} A & -B \\ B & A \end{pmatrix} \in K,$$

where $U(n)$ denotes the unitary group of degree n . We identify $\mathfrak{p}$ (resp. K) with T_n (resp. $U(n)$) through the map Ψ (resp. δ). We consider the action of $U(n)$ on T_n defined by

$$(2.6) \quad h \cdot Z = hZ^t h, \quad h \in U(n), \quad Z \in T_n.$$

Then the adjoint action (2.3) of K on $\mathfrak{p}$ is compatible with the action (2.6) of $U(n)$ on T_n through the map Ψ . Precisely for any $k \in K$ and $\omega \in \mathfrak{p}$, we get

$$(2.7) \quad \Psi(k\omega^t k) = \delta(k)\Psi(\omega)^t \delta(k).$$

The action (2.6) induces the action of $U(n)$ on the polynomial algebra $\text{Pol}(T_n)$ and the symmetric algebra $S(T_n)$ respectively. We denote by $\text{Pol}(T_n)^{U(n)}$ (resp. $S(T_n)^{U(n)}$) the subalgebra of $\text{Pol}(T_n)$ (resp. $S(T_n)$) consisting of $U(n)$ -invariants. The following inner product $(\ , \)$ on T_n defined by

$$(Z, W) = \text{tr}(Z \overline{W}), \quad Z, W \in T_n$$

gives an isomorphism as vector spaces

$$(2.8) \quad T_n \cong T_n^*, \quad Z \mapsto f_Z, \quad Z \in T_n,$$

where T_n^* denotes the dual space of T_n and f_Z is the linear functional on T_n defined by

$$f_Z(W) = (W, Z), \quad W \in T_n.$$

It is known that there is a canonical linear bijection of $S(T_n)^{U(n)}$ onto the algebra $\mathbb{D}(\mathbb{H}_n)$ of differential operators on $\mathbb{H}_n$ invariant under the action (1.1) of G . Identifying T_n with T_n^* by the above isomorphism (2.8), we get a canonical linear bijection

$$(2.9) \quad \Phi : \text{Pol}(T_n)^{U(n)} \longrightarrow \mathbb{D}(\mathbb{H}_n)$$

of $\text{Pol}(T_n)^{U(n)}$ onto $\mathbb{D}(\mathbb{H}_n)$. The map Φ is described explicitly as follows. Similarly the action (2.3) induces the action of K on the polynomial algebra $\text{Pol}(\mathfrak{p})$ and the symmetric algebra $S(\mathfrak{p})$ respectively. Through the map Ψ , the subalgebra $\text{Pol}(\mathfrak{p})^K$ of $\text{Pol}(\mathfrak{p})$ consisting of K -invariants is isomorphic to $\text{Pol}(T_n)^{U(n)}$. We put $N = n(n+1)$. Let $\{\xi_\alpha \mid 1 \leq \alpha \leq N\}$ be a basis of $\mathfrak{p}$. If $P \in \text{Pol}(\mathfrak{p})^K$, then

$$(2.10) \quad (\Phi(P)f)(gK) = \left[P \left(\frac{\partial}{\partial t_\alpha} \right) f \left(g \exp \left(\sum_{\alpha=1}^N t_\alpha \xi_\alpha \right) K \right) \right]_{(t_\alpha)=0},$$

where $f \in C^\infty(\mathbb{H}_n)$. We refer to [6, 7] for more detail. In general, it is hard to express $\Phi(P)$ explicitly for a polynomial $P \in \text{Pol}(\mathfrak{p})^K$.

According to the work of Harish-Chandra [4, 5], the algebra $\mathbb{D}(\mathbb{H}_n)$ is generated by n algebraically independent generators and is isomorphic to the commutative ring

$\mathbb{C}[x_1, \dots, x_n]$ with n indeterminates. We note that n is the real rank of G . Let $\mathfrak{g}_{\mathbb{C}}$ be the complexification of $\mathfrak{g}$. It is known that $\mathbb{D}(\mathbb{H}_n)$ is isomorphic to the center of the universal enveloping algebra of $\mathfrak{g}_{\mathbb{C}}$.

Using a classical invariant theory (cf. [8, 15], we can show that $\text{Pol}(T_n)^{U(n)}$ is generated by the following algebraically independent polynomials

$$(2.11) \quad q_j(Z) = \text{tr} \left((Z \overline{Z})^j \right), \quad j = 1, 2, \dots, n.$$

For each j with $1 \leq j \leq n$, the image $\Phi(q_j)$ of q_j is an invariant differential operator on $\mathbb{H}_n$ of degree $2j$. The algebra $\mathbb{D}(\mathbb{H}_n)$ is generated by n algebraically independent generators $\Phi(q_1), \Phi(q_2), \dots, \Phi(q_n)$. In particular,

$$(2.12) \quad \Phi(q_1) = c_1 \text{tr} \left(Y \left(Y \frac{\partial}{\partial \overline{\Omega}} \right) \frac{\partial}{\partial \Omega} \right) \quad \text{for some constant } c_1.$$

We observe that if we take $Z = X + iY$ with real X, Y , then $q_1(Z) = q_1(X, Y) = \text{tr}(X^2 + Y^2)$ and

$$q_2(Z) = q_2(X, Y) = \text{tr} \left((X^2 + Y^2)^2 + 2X(XY - YX)Y \right).$$

It is a natural question to express the images $\Phi(q_j)$ explicitly for $j = 2, 3, \dots, n$. We hope that the images $\Phi(q_j)$ for $j = 2, 3, \dots, n$ are expressed in the form of the *trace* as $\Phi(q_1)$.

H. Maass [10] found algebraically independent generators $H_1, H_2, \dots, H_n$ of $\mathbb{D}(\mathbb{H}_n)$. We will describe $H_1, H_2, \dots, H_n$ explicitly. For $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in Sp(n, \mathbb{R})$ and $\Omega = X + iY \in \mathbb{H}_n$ with real X, Y , we set

$$\Omega_* = M \cdot \Omega = X_* + iY_* \quad \text{with } X_*, Y_* \text{ real.}$$

We set

$$\begin{aligned} K &= (\Omega - \overline{\Omega}) \frac{\partial}{\partial \Omega} = 2iY \frac{\partial}{\partial \Omega}, \\ \Lambda &= (\Omega - \overline{\Omega}) \frac{\partial}{\partial \overline{\Omega}} = 2iY \frac{\partial}{\partial \overline{\Omega}}, \\ K_* &= (\Omega_* - \overline{\Omega}_*) \frac{\partial}{\partial \Omega_*} = 2iY_* \frac{\partial}{\partial \Omega_*}, \\ \Lambda_* &= (\Omega_* - \overline{\Omega}_*) \frac{\partial}{\partial \overline{\Omega}_*} = 2iY_* \frac{\partial}{\partial \overline{\Omega}_*}. \end{aligned}$$

Then it is easily seen that

$$(2.13) \quad K_* = {}^t(C\overline{\Omega} + D)^{-1} {}^t \{ (C\Omega + D) {}^t K \},$$

$$(2.14) \quad \Lambda_* = {}^t(C\Omega + D)^{-1} {}^t \{ (C\overline{\Omega} + D) {}^t \Lambda \}$$

and

$$(2.15) \quad {}^t\{(C\overline{\Omega} + D) {}^t\Lambda\} = \Lambda {}^t(C\overline{\Omega} + D) - \frac{n+1}{2} (\Omega - \overline{\Omega}) {}^tC.$$

Using Formulas (2.13), (2.14) and (2.15), we can show that

$$(2.16) \quad \Lambda_* K_* + \frac{n+1}{2} K_* = {}^t(C\Omega + D)^{-1} \left\{ (C\Omega + D) \left(\Lambda K + \frac{n+1}{2} K \right) \right\}.$$

Therefore we get

$$(2.17) \quad \text{tr} \left(\Lambda_* K_* + \frac{n+1}{2} K_* \right) = \text{tr} \left(\Lambda K + \frac{n+1}{2} K \right).$$

We set

$$(2.18) \quad A^{(1)} = \Lambda K + \frac{n+1}{2} K.$$

We define $A^{(j)}$ ($j = 2, 3, \dots, n$) recursively by

$$(2.19) \quad \begin{aligned} A^{(j)} &= A^{(1)} A^{(j-1)} - \frac{n+1}{2} \Lambda A^{(j-1)} + \frac{1}{2} \Lambda \sigma(A^{(j-1)}) \\ &\quad + \frac{1}{2} (\Omega - \overline{\Omega}) {}^t\left\{ (\Omega - \overline{\Omega})^{-1} {}^t({}^t\Lambda A^{(j-1)}) \right\}. \end{aligned}$$

We set

$$(2.20) \quad H_j = \text{tr}(A^{(j)}), \quad j = 1, 2, \dots, n.$$

As mentioned before, Maass proved that $H_1, H_2, \dots, H_n$ are algebraically independent generators $H_1, H_2, \dots, H_n$ of $\mathbb{D}(\mathbb{H}_n)$.

In fact, we see that

$$(2.21) \quad -H_1 = \Delta_{n;1} = 4 \text{tr} \left(Y {}^t \left(Y \frac{\partial}{\partial \overline{\Omega}} \right) \frac{\partial}{\partial \Omega} \right).$$

is the Laplacian for the invariant metric $ds_{n;1}^2$ on $\mathbb{H}_n$.

Conjecture. For $j = 2, 3, \dots, n$, $\Phi(q_j) = c_j H_j$ for a suitable constant c_j .

Example 2.1. We consider the case $n = 1$. The algebra $\text{Pol}(T_1)^{U(1)}$ is generated by the polynomial

$$q(z) = z \overline{z}, \quad z \in \mathbb{C}.$$

Using Formula (2.10), we get

$$\Phi(q) = 4y^2 \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right).$$

Therefore $\mathbb{D}(\mathbb{H}_1) = \mathbb{C}[\Phi(q)] = \mathbb{C}[H_1]$.

Example 2.2. We consider the case $n = 2$. The algebra $\text{Pol}(T_2)^{U(2)}$ is generated by the polynomial

$$q_1(Z) = \text{tr}(Z \overline{Z}), \quad q_2(Z) = \text{tr}\left((Z \overline{Z})^2\right), \quad Z \in T_2.$$

Using Formula (2.10), we may express $\Phi(q_1)$ and $\Phi(q_2)$ explicitly. $\Phi(q_1)$ is expressed by Formula (2.12). The computation of $\Phi(q_2)$ might be quite tedious. We leave the detail to the reader. In this case, $\Phi(q_2)$ was essentially computed in [2], Proposition 6. Therefore

$$\mathbb{D}(\mathbb{H}_2) = \mathbb{C}[\Phi(q_1), \Phi(q_2)] = \mathbb{C}[H_1, H_2].$$

They computed the center of the universal enveloping algebra $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$.

G. Shimura [12] found canonically defined algebraically independent generators of $\mathbb{D}(\mathbb{H}_n)$. We will describe his way of constructing those generators roughly. Let $K_{\mathbb{C}}, \mathfrak{g}_{\mathbb{C}}, \mathfrak{k}_{\mathbb{C}}, \mathfrak{p}_{\mathbb{C}}, \dots$ denote the complexification of $K, \mathfrak{g}, \mathfrak{k}, \mathfrak{p}, \dots$ respectively. Then we have the Cartan decomposition

$$\mathfrak{g}_{\mathbb{C}} = \mathfrak{k}_{\mathbb{C}} + \mathfrak{p}_{\mathbb{C}}, \quad \mathfrak{p}_{\mathbb{C}} = \mathfrak{p}_{\mathbb{C}}^+ + \mathfrak{p}_{\mathbb{C}}^-$$

with the properties

$$[\mathfrak{k}_{\mathbb{C}}, \mathfrak{p}_{\mathbb{C}}^{\pm}] \subset \mathfrak{p}_{\mathbb{C}}^{\pm}, \quad [\mathfrak{p}_{\mathbb{C}}^+, \mathfrak{p}_{\mathbb{C}}^+] = [\mathfrak{p}_{\mathbb{C}}^-, \mathfrak{p}_{\mathbb{C}}^-] = \{0\}, \quad [\mathfrak{p}_{\mathbb{C}}^+, \mathfrak{p}_{\mathbb{C}}^-] = \mathfrak{k}_{\mathbb{C}},$$

where

$$\begin{aligned} \mathfrak{g}_{\mathbb{C}} &= \left\{ \begin{pmatrix} X_1 & X_2 \\ X_3 & -{}^t X_1 \end{pmatrix} \mid X_1, X_2, X_3 \in \mathbb{C}^{(n,n)}, X_2 = {}^t X_2, X_3 = {}^t X_3 \right\}, \\ \mathfrak{k}_{\mathbb{C}} &= \left\{ \begin{pmatrix} A & -B \\ B & A \end{pmatrix} \in \mathbb{C}^{(2n,2n)} \mid {}^t A + A = 0, B = {}^t B \right\}, \\ \mathfrak{p}_{\mathbb{C}} &= \left\{ \begin{pmatrix} X & Y \\ Y & -X \end{pmatrix} \in \mathbb{C}^{(2n,2n)} \mid X = {}^t X, Y = {}^t Y \right\}, \\ \mathfrak{p}_{\mathbb{C}}^+ &= \left\{ \begin{pmatrix} Z & iZ \\ iZ & -Z \end{pmatrix} \in \mathbb{C}^{(2n,2n)} \mid Z = {}^t Z \in \mathbb{C}^{(n,n)} \right\}, \\ \mathfrak{p}_{\mathbb{C}}^- &= \left\{ \begin{pmatrix} Z & -iZ \\ -iZ & -Z \end{pmatrix} \in \mathbb{C}^{(2n,2n)} \mid Z = {}^t Z \in \mathbb{C}^{(n,n)} \right\}. \end{aligned}$$

For a complex vector space W and a nonnegative integer r , we denote by $\text{Pol}_r(W)$ the vector space of complex-valued homogeneous polynomial functions on W of degree r . We put

$$\text{Pol}^r(W) := \sum_{s=0}^r \text{Pol}_s(W).$$

$\text{ML}_r(W)$ denotes the vector space of all $\mathbb{C}$ -multilinear maps of $W \times \dots \times W$ (r copies) into $\mathbb{C}$. An element Q of $\text{ML}_r(W)$ is called *symmetric* if

$$Q(x_1, \dots, x_r) = Q(x_{\pi(1)}, \dots, x_{\pi(r)})$$

for each permutation π of $\{1, 2, \dots, r\}$. Given $P \in \text{Pol}_r(W)$, there is a unique element symmetric element P_* of $\text{Ml}_r(W)$ such that

$$(2.22) \quad P(x) = P_*(x, \dots, x) \quad \text{for all } x \in W.$$

Moreover the map $P \mapsto P_*$ is a $\mathbb{C}$ -linear bijection of $\text{Pol}_r(W)$ onto the set of all symmetric elements of $\text{Ml}_r(W)$. We let $S_r(W)$ denote the subspace consisting of all homogeneous elements of degree r in the symmetric algebra $S(W)$. We note that $\text{Pol}_r(W)$ and $S_r(W)$ are dual to each other with respect to the pairing

$$(2.23) \quad \langle \alpha, x_1 \cdots x_r \rangle = \alpha_*(x_1, \dots, x_r) \quad (x_i \in W, \alpha \in \text{Pol}_r(W)).$$

Let $\mathfrak{p}_{\mathbb{C}}^*$ be the dual space of $\mathfrak{p}_{\mathbb{C}}$, that is, $\mathfrak{p}_{\mathbb{C}}^* = \text{Pol}_1(\mathfrak{p}_{\mathbb{C}})$. Let $\{X_1, \dots, X_N\}$ be a basis of $\mathfrak{p}_{\mathbb{C}}$ and $\{Y_1, \dots, Y_N\}$ be the basis of $\mathfrak{p}_{\mathbb{C}}^*$ dual to $\{X_\nu\}$, where $N = n(n+1)$. We note that $\text{Pol}_r(\mathfrak{p}_{\mathbb{C}})$ and $\text{Pol}_r(\mathfrak{p}_{\mathbb{C}}^*)$ are dual to each other with respect to the pairing

$$(2.24) \quad \langle \alpha, \beta \rangle = \sum \alpha_*(X_{i_1}, \dots, X_{i_r}) \beta_*(Y_{i_1}, \dots, Y_{i_r}),$$

where $\alpha \in \text{Pol}_r(\mathfrak{p}_{\mathbb{C}})$, $\beta \in \text{Pol}_r(\mathfrak{p}_{\mathbb{C}}^*)$ and $(i_1, \dots, i_r)$ runs over $\{1, \dots, N\}^r$. Let $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$ be the universal enveloping algebra of $\mathfrak{g}_{\mathbb{C}}$ and $\mathcal{U}^p(\mathfrak{g}_{\mathbb{C}})$ its subspace spanned by the elements of the form $V_1 \cdots V_s$ with $V_i \in \mathfrak{g}_{\mathbb{C}}$ and $s \leq p$. We recall that there is a $\mathbb{C}$ -linear bijection ψ of the symmetric algebra $S(\mathfrak{g}_{\mathbb{C}})$ of $\mathfrak{g}_{\mathbb{C}}$ onto $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$ which is characterized by the property that $\psi(X^r) = X^r$ for all $X \in \mathfrak{g}_{\mathbb{C}}$. For each $\alpha \in \text{Pol}_r(\mathfrak{p}_{\mathbb{C}}^*)$ we define an element $\omega(\alpha)$ of $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$ by

$$(2.25) \quad \omega(\alpha) := \sum \alpha_*(Y_{i_1}, \dots, Y_{i_r}) X_{i_1} \cdots X_{i_r},$$

where $(i_1, \dots, i_r)$ runs over $\{1, \dots, N\}^r$. If $Y \in \mathfrak{p}_{\mathbb{C}}$, then Y^r as an element of $\text{Pol}_r(\mathfrak{p}_{\mathbb{C}}^*)$ is defined by

$$Y^r(u) = Y(u)^r \quad \text{for all } u \in \mathfrak{p}_{\mathbb{C}}^*.$$

Hence $(Y^r)_*(u_1, \dots, u_r) = Y(u_1) \cdots Y(u_r)$. According to (2.25), we see that if $\alpha(\sum t_i Y_i) = P(t_1, \dots, t_N)$ for $t_i \in \mathbb{C}$ with a polynomial P , then

$$(2.26) \quad \omega(\alpha) = \psi(P(X_1, \dots, X_N)).$$

Thus ω is a $\mathbb{C}$ -linear injection of $\text{Pol}(\mathfrak{p}_{\mathbb{C}}^*)$ into $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$ independent of the choice of a basis. We observe that $\omega(\text{Pol}_r(\mathfrak{p}_{\mathbb{C}}^*)) = \psi(S_r(\mathfrak{p}_{\mathbb{C}}))$. It is a well-known fact that if $\alpha_1, \dots, \alpha_m \in \text{Pol}_r(\mathfrak{p}_{\mathbb{C}}^*)$, then

$$(2.27) \quad \omega(\alpha_1 \cdots \alpha_m) - \omega(\alpha_1) \cdots \omega(\alpha_m) \in \mathcal{U}^{r-1}(\mathfrak{g}_{\mathbb{C}}).$$

We have a canonical pairing

$$(2.28) \quad \langle \cdot, \cdot \rangle : \text{Pol}_r(\mathfrak{p}_{\mathbb{C}}^+) \times \text{Pol}_r(\mathfrak{p}_{\mathbb{C}}^-) \longrightarrow \mathbb{C}$$

defined by

$$(2.29) \quad \langle f, g \rangle = \sum f_*(\tilde{X}_{i_1}, \dots, \tilde{X}_{i_r}) g_*(\tilde{Y}_{i_1}, \dots, \tilde{Y}_{i_r}),$$

where f_* (resp. g_*) are the unique symmetric elements of $\text{Ml}_r(\mathfrak{p}_{\mathbb{C}}^+)$ (resp. $\text{Ml}_r(\mathfrak{p}_{\mathbb{C}}^-)$), and $\{\tilde{X}_1, \dots, \tilde{X}_{\tilde{N}}\}$ and $\{\tilde{Y}_1, \dots, \tilde{Y}_{\tilde{N}}\}$ are dual bases of $\mathfrak{p}_{\mathbb{C}}^+$ and $\mathfrak{p}_{\mathbb{C}}^-$ with respect to

the Killing form $B(X, Y) = 2(n+1) \operatorname{tr}(XY)$, $\tilde{N} = \frac{n(n+1)}{2}$, and $(i_1, \dots, i_r)$ runs over $\{1, \dots, \tilde{N}\}^r$.

The adjoint representation of $K_{\mathbb{C}}$ on $\mathfrak{p}_{\mathbb{C}}^{\pm}$ induces the representation of $K_{\mathbb{C}}$ on $\operatorname{Pol}_r(\mathfrak{p}_{\mathbb{C}}^{\pm})$. Given a $K_{\mathbb{C}}$ -irreducible subspace Z of $\operatorname{Pol}_r(\mathfrak{p}_{\mathbb{C}}^+)$, we can find a unique $K_{\mathbb{C}}$ -irreducible subspace W of $\operatorname{Pol}_r(\mathfrak{p}_{\mathbb{C}}^-)$ such that $\operatorname{Pol}_r(\mathfrak{p}_{\mathbb{C}}^-)$ is the direct sum of W and the annihilator of Z . Then Z and W are dual with respect to the pairing (2.28). Take bases $\{\zeta_1, \dots, \zeta_{\kappa}\}$ of Z and $\{\xi_1, \dots, \xi_{\kappa}\}$ of W that are dual to each other. We set

$$(2.30) \quad f_Z(x, y) = \sum_{\nu=1}^{\kappa} \zeta_{\nu}(x) \xi_{\nu}(y) \quad (x \in \mathfrak{p}_{\mathbb{C}}^+, y \in \mathfrak{p}_{\mathbb{C}}^-).$$

It is easily seen that f_Z belongs to $\operatorname{Pol}_{2r}(\mathfrak{p}_{\mathbb{C}})^K$ and is independent of the choice of dual bases $\{\zeta_{\nu}\}$ and $\{\xi_{\nu}\}$. Shimura [12] proved that there exists a canonically defined set $\{Z_1, \dots, Z_n\}$ with a $K_{\mathbb{C}}$ -irreducible subspace Z_r of $\operatorname{Pol}_r(\mathfrak{p}_{\mathbb{C}}^+)$ ($1 \leq r \leq n$) such that $f_{Z_1}, \dots, f_{Z_n}$ are algebraically independent generators of $\operatorname{Pol}(\mathfrak{p}_{\mathbb{C}})^K$. We can identify $\mathfrak{p}_{\mathbb{C}}^+$ with T_n . We can take Z_r as the subspace of $\operatorname{Pol}_r(T_n)$ spanned by the functions $f_{a,r}(Z) = \det_r({}^t a Z a)$ for all $a \in GL(n, \mathbb{C})$, where $\det_r(x)$ denotes the determinant of the upper left $r \times r$ submatrix of x . For every $f \in \operatorname{Pol}(\mathfrak{p}_{\mathbb{C}})^K$, we let $\Omega(f)$ denote the element of $\mathbb{D}(\mathbb{H}_n)$ represented by $\omega(f)$. Then $\mathbb{D}(\mathbb{H}_n)$ is the polynomial ring $\mathbb{C}[\omega(f_{Z_1}), \dots, \omega(f_{Z_n})]$ generated by n algebraically independent elements $\omega(f_{Z_1}), \dots, \omega(f_{Z_n})$.

3. Invariant differential operators on $\mathbb{H}_{n,m}$

The stabilizer K^J of G^J at $(iI_n, 0)$ is given by

$$K^J = \left\{ (k, (0, 0; \kappa)) \mid k \in K, \kappa = {}^t \kappa \in \mathbb{R}^{(m,m)} \right\}.$$

Therefore $\mathbb{H}_{n,m} \cong G^J / K^J$ is a homogeneous space of *non-reductive type*. The Lie algebra $\mathfrak{g}^J$ of G^J has a decomposition

$$\mathfrak{g}^J = \mathfrak{k}^J + \mathfrak{p}^J,$$

where

$$\begin{aligned} \mathfrak{k}^J &= \left\{ (X, (0, 0, \kappa)) \mid X \in \mathfrak{k}, \kappa = {}^t \kappa \in \mathbb{R}^{(m,m)} \right\}, \\ \mathfrak{p}^J &= \left\{ (Y, (P, Q, 0)) \mid Y \in \mathfrak{p}, P, Q \in \mathbb{R}^{(m,n)} \right\}. \end{aligned}$$

Thus the tangent space of the homogeneous space $\mathbb{H}_{n,m}$ at $(iI_n, 0)$ is identified with $\mathfrak{p}^J$. We can see that

$$[\mathfrak{k}^J, \mathfrak{k}^J] \subset \mathfrak{k}^J, \quad [\mathfrak{k}^J, \mathfrak{p}^J] \subset \mathfrak{p}^J.$$

The adjoint action of K^J on $\mathfrak{p}^J$ induces the action of K on $\mathfrak{p}^J$ defined by

$$(3.1) \quad k \cdot Z = k Z {}^t k, \quad k \in K, \quad Z \in \mathfrak{p}^J.$$

Identifying $\mathbb{R}^{(m,n)} \times \mathbb{R}^{(m,n)}$ with $\mathbb{C}^{(m,n)}$, we can identify $\mathfrak{p}^J$ with $T_n \times \mathbb{C}^{(m,n)}$. For brevity, we write $T_{n,m} = T_n \times \mathbb{C}^{(m,n)}$. Through this identification the action (3.1) is compatible with the action of $U(n)$ on $T_{n,m}$ defined by

$$(3.2) \quad h \cdot (\omega, z) = (h \omega {}^t h, z {}^t h),$$

where $h \in K$, $\omega \in T_n$ and $z \in \mathbb{C}^{(m,n)}$. Here we regard the complex vector space $T_{n,m}$ as a real vector space.

We now describe the algebra $\mathbb{D}(\mathbb{H}_{n,m})$ of all differential operators on $\mathbb{H}_{n,m}$ invariant under the action (1.2) of G^J . The action (3.2) induces the action of $U(n)$ on the polynomial algebra $\text{Pol}_{n,m} = \text{Pol}(T_{n,m})$. We denote by $\text{Pol}_{n,m}^{U(n)}$ the subalgebra of $\text{Pol}_{n,m}$ consisting of all $U(n)$ -invariants. Similarly the action (3.1) of K induces the action of K on the polynomial algebra $\text{Pol}(\mathfrak{p}^J)$. We see that through the identification of $\mathfrak{p}^J$ with $T_{n,m}$, the algebra $\text{Pol}(\mathfrak{p}^J)$ is isomorphic to $\text{Pol}_{n,m}$. The following $U(n)$ -invariant inner product $(\cdot, \cdot)_*$ of the complex vector space $T_{n,m}$ defined by

$$((\omega, z), (\omega', z'))_* = \text{tr}(\omega \overline{\omega'}) + \text{tr}(z \overline{{}^t z'}), \quad (\omega, z), (\omega', z') \in T_{n,m}$$

gives a canonical isomorphism

$$T_{n,m} \cong T_{n,m}^*, \quad (\omega, z) \mapsto f_{\omega, z}, \quad (\omega, z) \in T_{n,m},$$

where $f_{\omega, z}$ is the linear functional on $T_{n,m}$ defined by

$$f_{\omega, z}((\omega', z')) = ((\omega', z'), (\omega, z))_*, \quad (\omega', z') \in T_{n,m}.$$

It is known that one gets a canonical linear bijection of $S(T_{n,m})^{U(n)}$ onto $\mathbb{D}(\mathbb{H}_{n,m})$. Identifying $T_{n,m}$ with $T_{n,m}^*$ by the above isomorphism, one gets a natural linear bijection

$$\Theta : \text{Pol}_{n,m}^{U(n)} \longrightarrow \mathbb{D}(\mathbb{H}_{n,m})$$

of $\text{Pol}_{n,m}^{U(n)}$ onto $\mathbb{D}(\mathbb{H}_{n,m})$. The map Θ is described explicitly as follows. We put $N_* = n(n+1) + 2mn$. Let $\{\eta_\alpha \mid 1 \leq \alpha \leq N_*\}$ be a basis of $\mathfrak{p}^J$. If $P \in \text{Pol}(\mathfrak{p}^J)^K = \text{Pol}_{n,m}^{U(n)}$, then

$$(3.3) \quad (\Theta(P)f)(gK) = \left[P \left(\frac{\partial}{\partial t_\alpha} \right) f \left(g \exp \left(\sum_{\alpha=1}^{N_*} t_\alpha \eta_\alpha \right) K \right) \right]_{(t_\alpha)=0},$$

where $f \in C^\infty(\mathbb{H}_{n,m})$. In general, it is hard to express $\Theta(P)$ explicitly for a polynomial $P \in \text{Pol}(\mathfrak{p}^J)^K$. We refer to [7], p. 287.

We present the following *basic* K -invariant polynomials in $\text{Pol}_{n,m}^{U(n)}$.

$$(3.4) \quad p_j(\omega, z) = \text{tr}((\omega \bar{\omega})^j), \quad 1 \leq j \leq n,$$

$$(3.5) \quad \psi_k^{(1)}(\omega, z) = (z^t \bar{z})_{kk}, \quad 1 \leq k \leq m,$$

$$(3.6) \quad \psi_{kp}^{(2)}(\omega, z) = \text{Re}(z^t \bar{z})_{kp}, \quad 1 \leq k < p \leq m,$$

$$(3.7) \quad \psi_{kp}^{(3)}(\omega, z) = \text{Im}(z^t \bar{z})_{kp}, \quad 1 \leq k < p \leq m,$$

$$(3.8) \quad f_{kp}^{(1)}(\omega, z) = \text{Re}(z \bar{\omega}^t z)_{kp}, \quad 1 \leq k \leq p \leq m,$$

$$(3.9) \quad f_{kp}^{(2)}(\omega, z) = \text{Im}(z \bar{\omega}^t z)_{kp}, \quad 1 \leq k \leq p \leq m,$$

where $\omega \in T_n$ and $z \in \mathbb{C}^{(m,n)}$.

We present some interesting K -invariants. For an $m \times m$ matrix S , we define the following invariant polynomials in $\text{Pol}_{n,m}^{U(n)}$:

$$(3.10) \quad m_{j;S}^{(1)}(\omega, z) = \text{Re} \left(\text{tr}(\omega \bar{\omega} + {}^t z S \bar{z})^j \right), \quad 1 \leq j \leq n,$$

$$(3.11) \quad m_{j;S}^{(2)}(\omega, z) = \text{Im} \left(\text{tr}(\omega \bar{\omega} + {}^t z S \bar{z})^j \right), \quad 1 \leq j \leq n,$$

$$(3.12) \quad q_{k;S}^{(1)}(\omega, z) = \text{Re} \left(\text{tr}(({}^t z S \bar{z})^k) \right), \quad 1 \leq k \leq m,$$

$$(3.13) \quad q_{k;S}^{(2)}(\omega, z) = \text{Im} \left(\text{tr}(({}^t z S \bar{z})^k) \right), \quad 1 \leq k \leq m,$$

$$(3.14) \quad \theta_{i,k,j;S}^{(1)}(\omega, z) = \text{Re} \left(\text{tr}((\omega \bar{\omega})^i ({}^t z S \bar{z})^k (\omega \bar{\omega} + {}^t z S \bar{z})^j) \right),$$

$$(3.15) \quad \theta_{i,k,j;S}^{(2)}(\omega, z) = \text{Im} \left(\text{tr}((\omega \bar{\omega})^i ({}^t z S \bar{z})^k (\omega \bar{\omega} + {}^t z S \bar{z})^j) \right),$$

where $1 \leq i, j \leq n$ and $1 \leq k \leq m$.

We define the following K -invariant polynomials in $\text{Pol}_{n,m}^{U(n)}$.

$$(3.16) \quad r_{jk}^{(1)}(\omega, z) = \text{Re} \left(\text{tr}((\omega \bar{\omega})^j ({}^t z \bar{z})^k) \right), \quad 1 \leq j \leq n, \quad 1 \leq k \leq m,$$

$$(3.17) \quad r_{jk}^{(2)}(\omega, z) = \text{Im} \left(\text{tr}((\omega \bar{\omega})^j ({}^t z \bar{z})^k) \right), \quad 1 \leq j \leq n, \quad 1 \leq k \leq m.$$

There may be possible other new invariants. We think that at this moment it may be complicated and difficult to find the generators of $\text{Pol}_{n,m}^{U(n)}$.

We propose the following problems.

Problem 1. Find the generators of $\text{Pol}_{n,m}^{U(n)}$. Are there canonical ways to find algebraically independent generators of $\text{Pol}_{n,m}^{U(n)}$ like Shimura's way in the hermitian symmetric space case ?

Problem 2. Find an easy way to express the images of the above invariant polynomials under the map Θ explicitly.

For $g = (M, (\lambda, \mu; \kappa)) \in G^J$ with $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in Sp(n, \mathbb{R})$ and $(\Omega, Z) \in \mathbb{H}_{n,m}$, we set

$$\begin{aligned}\Omega_* &= M \cdot \Omega = X_* + i Y_*, \quad X_*, Y_* \text{ real}, \\ Z_* &= (Z + \lambda \Omega + \mu)(C\Omega + D)^{-1} = U_* + i V_*, \quad U_*, V_* \text{ real}.\end{aligned}$$

For a coordinate $(\Omega, Z) \in \mathbb{H}_{n,m}$ with $\Omega = (\omega_{\mu\nu})$ and $Z = (z_{kl})$, we put $d\Omega$, $d\bar{\Omega}$, $\frac{\partial}{\partial\Omega}$, $\frac{\partial}{\partial\bar{\Omega}}$ as before and set

$$\begin{aligned}Z &= U + iV, \quad U = (u_{kl}), \quad V = (v_{kl}) \text{ real}, \\ dZ &= (dz_{kl}), \quad d\bar{Z} = (d\bar{z}_{kl}), \\ \frac{\partial}{\partial Z} &= \begin{pmatrix} \frac{\partial}{\partial z_{11}} & \cdots & \frac{\partial}{\partial z_{m1}} \\ \vdots & \ddots & \vdots \\ \frac{\partial}{\partial z_{1n}} & \cdots & \frac{\partial}{\partial z_{mn}} \end{pmatrix}, \quad \frac{\partial}{\partial \bar{Z}} = \begin{pmatrix} \frac{\partial}{\partial \bar{z}_{11}} & \cdots & \frac{\partial}{\partial \bar{z}_{m1}} \\ \vdots & \ddots & \vdots \\ \frac{\partial}{\partial \bar{z}_{1n}} & \cdots & \frac{\partial}{\partial \bar{z}_{mn}} \end{pmatrix}.\end{aligned}$$

Then we can show that

$$(3.18) \quad d\Omega_* = {}^t(C\Omega + D)^{-1} d\Omega (C\Omega + D)^{-1},$$

$$(3.19) \quad dZ_* = dZ(C\Omega + D)^{-1} + \{\lambda - (Z + \lambda\Omega + \mu)(C\Omega + D)^{-1}C\} d\Omega (C\Omega + D)^{-1},$$

$$(3.20) \quad \frac{\partial}{\partial\Omega_*} = (C\Omega + D) \left\{ (C\Omega + D) \frac{\partial}{\partial\Omega} \right\} + (C\Omega + D) \left\{ (C {}^tZ + C {}^t\mu - D {}^t\lambda) {}^t\left(\frac{\partial}{\partial Z}\right) \right\}$$

and

$$(3.21) \quad \frac{\partial}{\partial Z_*} = (C\Omega + D) \frac{\partial}{\partial Z}.$$

From [10, p. 33] or [14, p. 128], we know that

$$(3.22) \quad Y_* = {}^t(C\bar{\Omega} + D)^{-1} Y (C\Omega + D)^{-1} = {}^t(C\Omega + D)^{-1} Y (C\bar{\Omega} + D)^{-1}.$$

Using Formulas (3.18), (3.19) and (3.22), the author [22] proved that for any two positive real numbers A and B ,

$$\begin{aligned}ds_{n,m;A,B}^2 &= A \operatorname{tr} \left(Y^{-1} d\Omega Y^{-1} d\bar{\Omega} \right) \\ &\quad + B \left\{ \operatorname{tr} \left(Y^{-1} {}^tV V Y^{-1} d\Omega Y^{-1} d\bar{\Omega} \right) + \operatorname{tr} \left(Y^{-1} {}^t(dZ) d\bar{Z} \right) \right. \\ &\quad \left. - \operatorname{tr} \left(V Y^{-1} d\Omega Y^{-1} {}^t(d\bar{Z}) \right) - \operatorname{tr} \left(V Y^{-1} d\bar{\Omega} Y^{-1} {}^t(dZ) \right) \right\}\end{aligned}$$

is a Riemannian metric on $\mathbb{H}_{n,m}$ which is invariant under the action (1.2) of G^J .

Using Formulas (3.20), (3.21) and (3.22), we can prove that the following differential operators $\mathbb{M}_1$ and $\mathbb{M}_2$ on $\mathbb{H}_{n,m}$ defined by

$$(3.23) \quad \mathbb{M}_1 = \text{tr} \left(Y \frac{\partial}{\partial Z} {}^t \left(\frac{\partial}{\partial \bar{Z}} \right) \right)$$

and

$$(3.24) \quad \begin{aligned} \mathbb{M}_2 = & \text{tr} \left(Y {}^t \left(Y \frac{\partial}{\partial \bar{\Omega}} \right) \frac{\partial}{\partial \Omega} \right) + \text{tr} \left(V Y^{-1} {}^t V \left(Y \frac{\partial}{\partial \bar{Z}} \right) \frac{\partial}{\partial Z} \right) \\ & + \text{tr} \left(V {}^t \left(Y \frac{\partial}{\partial \bar{\Omega}} \right) \frac{\partial}{\partial Z} \right) + \text{tr} \left({}^t V \left(Y \frac{\partial}{\partial \bar{Z}} \right) \frac{\partial}{\partial \Omega} \right) \end{aligned}$$

are invariant under the action (1.2) of G^J . The author [22] proved that for any two positive real numbers A and B , the following differential operator

$$(3.25) \quad \Delta_{n,m;A,B} = \frac{4}{A} \mathbb{M}_2 + \frac{4}{B} \mathbb{M}_1$$

is the Laplacian of the G^J -invariant Riemannian metric $ds_{n,m;A,B}^2$.

Let

$$\mathbb{D}_n = \{ W \in \mathbb{C}^{(n,n)} \mid W = {}^t W, I_n - \bar{W}W > 0 \}$$

be the generalized unit disk.

For brevity, we write $\mathbb{D}_{n,m} := \mathbb{D}_n \times \mathbb{C}^{(m,n)}$. For a coordinate $(W, \eta) \in \mathbb{D}_{n,m}$ with $W = (w_{\mu\nu}) \in \mathbb{D}_n$ and $\eta = (\eta_{kl}) \in \mathbb{C}^{(m,n)}$, we put

$$\begin{aligned} dW &= (dw_{\mu\nu}), & d\bar{W} &= (d\bar{w}_{\mu\nu}), \\ d\eta &= (d\eta_{kl}), & d\bar{\eta} &= (d\bar{\eta}_{kl}) \end{aligned}$$

and

$$\begin{aligned} \frac{\partial}{\partial W} &= \left(\frac{1 + \delta_{\mu\nu}}{2} \frac{\partial}{\partial w_{\mu\nu}} \right), & \frac{\partial}{\partial \bar{W}} &= \left(\frac{1 + \delta_{\mu\nu}}{2} \frac{\partial}{\partial \bar{w}_{\mu\nu}} \right), \\ \frac{\partial}{\partial \eta} &= \begin{pmatrix} \frac{\partial}{\partial \eta_{11}} & \cdots & \frac{\partial}{\partial \eta_{m1}} \\ \vdots & \ddots & \vdots \\ \frac{\partial}{\partial \eta_{1n}} & \cdots & \frac{\partial}{\partial \eta_{mn}} \end{pmatrix}, & \frac{\partial}{\partial \bar{\eta}} &= \begin{pmatrix} \frac{\partial}{\partial \bar{\eta}_{11}} & \cdots & \frac{\partial}{\partial \bar{\eta}_{m1}} \\ \vdots & \ddots & \vdots \\ \frac{\partial}{\partial \bar{\eta}_{1n}} & \cdots & \frac{\partial}{\partial \bar{\eta}_{mn}} \end{pmatrix}. \end{aligned}$$

We can identify an element $g = (M, (\lambda, \mu; \kappa))$ of G^J , $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in Sp(n, \mathbb{R})$ with the element

$$\begin{pmatrix} A & 0 & B & A {}^t \mu - B {}^t \lambda \\ \lambda & I_m & \mu & \kappa \\ C & 0 & D & C {}^t \mu - D {}^t \lambda \\ 0 & 0 & 0 & I_m \end{pmatrix}$$

of $Sp(m+n, \mathbb{R})$.

We set

$$T_* = \frac{1}{\sqrt{2}} \begin{pmatrix} I_{m+n} & I_{m+n} \\ iI_{m+n} & -iI_{m+n} \end{pmatrix}.$$

We now consider the group G_*^J defined by

$$G_*^J := T_*^{-1} G^J T_*.$$

If $g = (M, (\lambda, \mu; \kappa)) \in G^J$ with $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in Sp(n, \mathbb{R})$, then $T_*^{-1} g T_*$ is given by

$$(3.26) \quad T_*^{-1} g T_* = \begin{pmatrix} P_* & Q_* \\ \overline{Q}_* & \overline{P}_* \end{pmatrix},$$

where

$$P_* = \begin{pmatrix} P & \frac{1}{2} \{Q^t(\lambda + i\mu) - P^t(\lambda - i\mu)\} \\ \frac{1}{2}(\lambda + i\mu) & I_h + i\frac{\kappa}{2} \end{pmatrix},$$

$$Q_* = \begin{pmatrix} Q & \frac{1}{2} \{P^t(\lambda - i\mu) - Q^t(\lambda + i\mu)\} \\ \frac{1}{2}(\lambda - i\mu) & -i\frac{\kappa}{2} \end{pmatrix},$$

and P, Q are given by the formulas

$$(3.27) \quad P = \frac{1}{2} \{(A + D) + i(B - C)\}$$

and

$$(3.28) \quad Q = \frac{1}{2} \{(A - D) - i(B + C)\}.$$

From now on, we write

$$\left(\begin{pmatrix} P & Q \\ \overline{Q} & \overline{P} \end{pmatrix}, \left(\frac{1}{2}(\lambda + i\mu), \frac{1}{2}(\lambda - i\mu); -i\frac{\kappa}{2} \right) \right) := \left(\frac{P_*}{Q_*}, \frac{Q_*}{P_*} \right).$$

In other words, we have the relation

$$T_*^{-1} \left(\begin{pmatrix} A & B \\ C & D \end{pmatrix}, (\lambda, \mu; \kappa) \right) T_* = \left(\begin{pmatrix} P & Q \\ \overline{Q} & \overline{P} \end{pmatrix}, \left(\frac{1}{2}(\lambda + i\mu), \frac{1}{2}(\lambda - i\mu); -i\frac{\kappa}{2} \right) \right).$$

Let

$$H_{\mathbb{C}}^{(n,m)} := \{(\xi, \eta; \zeta) \mid \xi, \eta \in \mathbb{C}^{(m,n)}, \zeta \in \mathbb{C}^{(m,m)}, \zeta + \eta^t \xi \text{ symmetric}\}$$

be the complex Heisenberg group endowed with the following multiplication

$$(\xi, \eta; \zeta) \circ (\xi', \eta'; \zeta') := (\xi + \xi', \eta + \eta'; \zeta + \zeta' + \xi^t \eta' - \eta^t \xi').$$

We define the semidirect product

$$SL(2n, \mathbb{C}) \ltimes H_{\mathbb{C}}^{(n,m)}$$

endowed with the following multiplication

$$\begin{aligned} & \left(\begin{pmatrix} P & Q \\ R & S \end{pmatrix}, (\xi, \eta; \zeta) \right) \cdot \left(\begin{pmatrix} P' & Q' \\ R' & S' \end{pmatrix}, (\xi', \eta'; \zeta') \right) \\ &= \left(\begin{pmatrix} P & Q \\ R & S \end{pmatrix} \begin{pmatrix} P' & Q' \\ R' & S' \end{pmatrix}, (\tilde{\xi} + \xi', \tilde{\eta} + \eta'; \zeta + \zeta' + \tilde{\xi}^t \eta' - \tilde{\eta}^t \xi') \right), \end{aligned}$$

where $\tilde{\xi} = \xi P' + \eta R'$ and $\tilde{\eta} = \xi Q' + \eta S'$.

If we identify $H_{\mathbb{R}}^{(n,m)}$ with the subgroup

$$\{(\xi, \bar{\xi}; i\kappa) \mid \xi \in \mathbb{C}^{(m,n)}, \kappa \in \mathbb{R}^{(m,m)}\}$$

of $H_{\mathbb{C}}^{(n,m)}$, we have the following inclusion

$$G_*^J \subset SU(n, n) \ltimes H_{\mathbb{R}}^{(n,m)} \subset SL(2n, \mathbb{C}) \ltimes H_{\mathbb{C}}^{(n,m)}.$$

We define the mapping $\Theta : G^J \longrightarrow G_*^J$ by

$$(3.29) \quad \Theta \left(\begin{pmatrix} A & B \\ C & D \end{pmatrix}, (\lambda, \mu; \kappa) \right) := \left(\begin{pmatrix} P & Q \\ \bar{Q} & \bar{P} \end{pmatrix}, \left(\frac{1}{2}(\lambda + i\mu), \frac{1}{2}(\lambda - i\mu); -i\frac{\kappa}{2} \right) \right),$$

where P and Q are given by (3.27) and (3.28). We can see that if $g_1, g_2 \in G^J$, then $\Theta(g_1 g_2) = \Theta(g_1) \Theta(g_2)$.

According to [20, p. 250], G_*^J is of the Harish-Chandra type (cf. [11, p. 118]). Let

$$g_* = \left(\begin{pmatrix} P & Q \\ \bar{Q} & \bar{P} \end{pmatrix}, (\lambda, \mu; \kappa) \right)$$

be an element of G_*^J . Since the Harish-Chandra decomposition of an element $\begin{pmatrix} P & Q \\ R & S \end{pmatrix}$ in $SU(n, n)$ is given by

$$\begin{pmatrix} P & Q \\ R & S \end{pmatrix} = \begin{pmatrix} I_n & QS^{-1} \\ 0 & I_n \end{pmatrix} \begin{pmatrix} P - QS^{-1}R & 0 \\ 0 & S \end{pmatrix} \begin{pmatrix} I_n & 0 \\ S^{-1}R & I_n \end{pmatrix},$$

the P_*^+ -component of the following element

$$g_* \cdot \left(\begin{pmatrix} I_n & W \\ 0 & I_n \end{pmatrix}, (0, \eta; 0) \right), \quad W \in \mathbb{D}_n$$

of $SL(2n, \mathbb{C}) \ltimes H_{\mathbb{C}}^{(n,m)}$ is given by

$$(3.30) \quad \left(\begin{pmatrix} I_n & (PW + Q)(\bar{Q}W + \bar{P})^{-1} \\ 0 & I_n \end{pmatrix}, (0, (\eta + \lambda W + \mu)(\bar{Q}W + \bar{P})^{-1}; 0) \right).$$

We can identify $\mathbb{D}_{n,m}$ with the subset

$$\left\{ \left(\begin{pmatrix} I_n & W \\ 0 & I_n \end{pmatrix}, (0, \eta; 0) \right) \mid W \in \mathbb{D}_n, \eta \in \mathbb{C}^{(m,n)} \right\}$$

of the complexification of G_*^J . Indeed, $\mathbb{D}_{n,m}$ is embedded into P_*^+ given by

$$P_*^+ = \left\{ \left(\begin{pmatrix} I_n & W \\ 0 & I_n \end{pmatrix}, (0, \eta; 0) \right) \mid W = {}^t W \in \mathbb{C}^{(n,n)}, \eta \in \mathbb{C}^{(m,n)} \right\}.$$

This is a generalization of the Harish-Chandra embedding (cf. [11, p. 119]). Then we get the *natural transitive action* of G_*^J on $\mathbb{D}_{n,m}$ defined by

$$(3.31) \quad \left(\begin{pmatrix} P & Q \\ \overline{Q} & \overline{P} \end{pmatrix}, (\xi, \bar{\xi}; i\kappa) \right) \cdot (W, \eta) \\ = \left((PW + Q)(\overline{Q}W + \overline{P})^{-1}, (\eta + \xi W + \bar{\xi})(\overline{Q}W + \overline{P})^{-1} \right),$$

where $\begin{pmatrix} P & Q \\ \overline{Q} & \overline{P} \end{pmatrix} \in G_*$, $\xi \in \mathbb{C}^{(m,n)}$, $\kappa \in \mathbb{R}^{(m,m)}$ and $(W, \eta) \in \mathbb{D}_{n,m}$.

The author [23] proved that the action (1.2) of G^J on $\mathbb{H}_{n,m}$ is compatible with the action (3.31) of G_*^J on $\mathbb{D}_{n,m}$ through a *partial Cayley transform* $\Phi : \mathbb{D}_{n,m} \longrightarrow \mathbb{H}_{n,m}$ defined by

$$(3.32) \quad \Phi(W, \eta) := \left(i(I_n + W)(I_n - W)^{-1}, 2i\eta(I_n - W)^{-1} \right).$$

In other words, if $g_0 \in G^J$ and $(W, \eta) \in \mathbb{D}_{n,m}$,

$$(3.33) \quad g_0 \cdot \Phi(W, \eta) = \Phi(g_* \cdot (W, \eta)),$$

where $g_* = T_*^{-1}g_0T_*$. Φ is a biholomorphic mapping of $\mathbb{D}_{n,m}$ onto $\mathbb{H}_{n,m}$ which gives the partially bounded realization of $\mathbb{H}_{n,m}$ by $\mathbb{D}_{n,m}$. The inverse of Φ is

$$\Phi^{-1}(\Omega, Z) = \left((\Omega - iI_n)(\Omega + iI_n)^{-1}, Z(\Omega + iI_n)^{-1} \right).$$

For $(W, \eta) \in \mathbb{D}_{n,m}$, we write

$$(\Omega, Z) := \Phi(W, \eta).$$

Thus

$$(3.34) \quad \Omega = i(I_n + W)(I_n - W)^{-1}, \quad Z = 2i\eta(I_n - W)^{-1}.$$

Since

$$d(I_n - W)^{-1} = (I_n - W)^{-1}dW(I_n - W)^{-1}$$

and

$$I_n + (I_n + W)(I_n - W)^{-1} = 2(I_n - W)^{-1},$$

we get the following formulas from (3.34)

$$(3.35) \quad Y = \frac{1}{2i}(\Omega - \overline{\Omega}) = (I_n - W)^{-1}(I_n - W\overline{W})(I_n - \overline{W})^{-1},$$

$$(3.36) \quad V = \frac{1}{2i}(Z - \overline{Z}) = \eta(I_n - W)^{-1} + \overline{\eta}(I_n - \overline{W})^{-1},$$

$$(3.37) \quad d\Omega = 2i(I_n - W)^{-1}dW(I_n - W)^{-1},$$

$$(3.38) \quad dZ = 2i \left\{ d\eta + \eta(I_n - W)^{-1}dW \right\} (I_n - W)^{-1}.$$

Using Formulas (3.32), (3.34)-(3.38), we can prove that for any two positive real numbers A and B , the following metric $ds_{\mathbb{D}_{n,m};A,B}^2$ defined by

$$\begin{aligned}
ds_{\mathbb{D}_{n,m};A,B}^2 = & 4A \operatorname{tr} \left((I_n - W\overline{W})^{-1} dW (I_n - \overline{W}W)^{-1} d\overline{W} \right) \\
& + 4B \left\{ \operatorname{tr} \left((I_n - W\overline{W})^{-1} {}^t(d\eta) \beta \right) \right. \\
& + \operatorname{tr} \left((\eta\overline{W} - \overline{\eta})(I_n - W\overline{W})^{-1} dW (I_n - \overline{W}W)^{-1} {}^t(d\overline{\eta}) \right) \\
& + \operatorname{tr} \left((\overline{\eta}W - \eta)(I_n - \overline{W}W)^{-1} d\overline{W} (I_n - W\overline{W})^{-1} {}^t(d\eta) \right) \\
& - \operatorname{tr} \left((I_n - W\overline{W})^{-1} {}^t\eta \eta (I_n - \overline{W}W)^{-1} \overline{W} dW (I_n - \overline{W}W)^{-1} d\overline{W} \right) \\
& - \operatorname{tr} \left(W (I_n - \overline{W}W)^{-1} {}^t\overline{\eta} \overline{\eta} (I_n - W\overline{W})^{-1} dW (I_n - \overline{W}W)^{-1} d\overline{W} \right) \\
& + \operatorname{tr} \left((I_n - W\overline{W})^{-1} {}^t\eta \overline{\eta} (I_n - W\overline{W})^{-1} dW (I_n - \overline{W}W)^{-1} d\overline{W} \right) \\
& + \operatorname{tr} \left((I_n - \overline{W})^{-1} {}^t\overline{\eta} \eta \overline{W} (I_n - W\overline{W})^{-1} dW (I_n - \overline{W}W)^{-1} d\overline{W} \right) \\
& + \operatorname{tr} \left((I_n - \overline{W})^{-1} (I_n - W) (I_n - \overline{W}W)^{-1} {}^t\overline{\eta} \eta (I_n - \overline{W}W)^{-1} \right. \\
& \quad \times (I_n - \overline{W}) (I_n - W)^{-1} dW (I_n - \overline{W}W)^{-1} d\overline{W} \left. \right) \\
& \left. - \operatorname{tr} \left((I_n - W\overline{W})^{-1} (I_n - W) (I_n - \overline{W})^{-1} {}^t\overline{\eta} \eta (I_n - W)^{-1} \right. \right. \\
& \quad \left. \left. \times dW (I_n - \overline{W}W)^{-1} d\overline{W} \right) \right\}
\end{aligned}$$

is a Riemannian metric on $\mathbb{D}_{n,m}$ which is invariant under the action (3.31) of the Jacobi group G_*^J .

We note that if $n = m = 1$ and $A = B = 1$, we get

$$\begin{aligned}
\frac{1}{4} ds_{\mathbb{D}_{1,1};1,1}^2 = & \frac{dW d\overline{W}}{(1 - |W|^2)^2} + \frac{1}{(1 - |W|^2)} d\eta d\overline{\eta} \\
& + \frac{(1 + |W|^2)|\eta|^2 - \overline{W}\eta^2 - W\overline{\eta}^2}{(1 - |W|^2)^3} dW d\overline{W} \\
& + \frac{\eta\overline{W} - \overline{\eta}}{(1 - |W|^2)^2} dW d\overline{\eta} + \frac{\overline{\eta}W - \eta}{(1 - |W|^2)^2} d\overline{W} d\eta.
\end{aligned}$$

From the formulas (3.34), (3.37) and (3.38), we get

$$(3.39) \quad \frac{\partial}{\partial \Omega} = \frac{1}{2i} (I_n - W) \left[{}^t \left\{ (I_n - W) \frac{\partial}{\partial W} \right\} - {}^t \left\{ {}^t\eta \left(\frac{\partial}{\partial \eta} \right) \right\} \right]$$

and

$$(3.40) \quad \frac{\partial}{\partial Z} = \frac{1}{2i} (I_n - W) \frac{\partial}{\partial \eta}.$$

The following lemma is very useful for computing the invariant differential operators. H. Maass [9] observed the following useful fact.

Lemma 3.1. (a) Let A be an $m \times n$ matrix and B an $n \times l$ matrix. Assume that the entries of A commute with the entries of B . Then ${}^t(AB) = {}^tB {}^tA$.

(b) Let A , B and C be a $k \times l$, an $n \times m$ and an $m \times l$ matrix respectively. Assume that the entries of A commute with the entries of B . Then

$${}^t(A {}^t(BC)) = B {}^t(A {}^tC).$$

Proof. The proof follows immediately from the direct computation. $\square$

Using Formulas (3.34)-(3.36), (3.39), (3.40) and Lemma 3.1, we can prove that the following differential operators $\mathbb{S}_1$ and $\mathbb{S}_2$ on $\mathbb{D}_{n,m}$ defined by

$$\mathbb{S}_1 = \sigma \left((I_n - \overline{W}W) \frac{\partial}{\partial \eta} \left(\frac{\partial}{\partial \overline{\eta}} \right) \right)$$

and

$$\begin{aligned} \mathbb{S}_2 = & \operatorname{tr} \left((I_n - W\overline{W}) {}^t \left((I_n - W\overline{W}) \frac{\partial}{\partial \overline{W}} \right) \frac{\partial}{\partial W} \right) \\ & + \operatorname{tr} \left({}^t(\eta - \overline{\eta}W) \left(\frac{\partial}{\partial \overline{\eta}} \right) (I_n - \overline{W}W) \frac{\partial}{\partial W} \right) \\ & + \operatorname{tr} \left((\overline{\eta} - \eta\overline{W}) {}^t \left((I_n - W\overline{W}) \frac{\partial}{\partial \overline{W}} \right) \frac{\partial}{\partial \eta} \right) \\ & - \operatorname{tr} \left(\eta\overline{W}(I_n - W\overline{W})^{-1} {}^t \left(\frac{\partial}{\partial \overline{\eta}} \right) (I_n - \overline{W}W) \frac{\partial}{\partial \eta} \right) \\ & - \operatorname{tr} \left(\overline{\eta}W(I_n - \overline{W}W)^{-1} {}^t \left(\frac{\partial}{\partial \overline{\eta}} \right) (I_n - \overline{W}W) \frac{\partial}{\partial \eta} \right) \\ & + \operatorname{tr} \left(\overline{\eta}(I_n - W\overline{W})^{-1} {}^t \left(\frac{\partial}{\partial \overline{\eta}} \right) (I_n - \overline{W}W) \frac{\partial}{\partial \eta} \right) \\ & + \operatorname{tr} \left(\eta\overline{W}W(I_n - \overline{W}W)^{-1} {}^t \left(\frac{\partial}{\partial \overline{\eta}} \right) (I_n - \overline{W}W) \frac{\partial}{\partial \eta} \right) \end{aligned}$$

are invariant under the action (3.31) of G_*^J . We can prove that

$$(3.41) \quad \Delta_{\mathbb{D}_{n,m};A,B} := \frac{1}{A} \mathbb{S}_2 + \frac{1}{B} \mathbb{S}_1$$

is the Laplacian of the invariant metric $ds_{\mathbb{D}_{n,m};A,B}^2$ on $\mathbb{D}_{n,m}$ (cf. [25]).

Using (3.21), (3.22) and Lemma 3.1, we can see that the following differential operator $\mathbb{K}$ on $\mathbb{H}_{n,m}$ of degree $2n$ defined by

$$(3.42) \quad \mathbb{K} = \det(Y) \det \left(\frac{\partial}{\partial Z} {}^t \left(\frac{\partial}{\partial \bar{Z}} \right) \right)$$

is invariant under the action (1.2) of G^J . In fact, if f is a C^∞ function on $\mathbb{H}_{n,m}$,

$$\begin{aligned} \mathbb{K}_* f &= \det(Y) |\det(C\Omega + D)|^{-2} \det \left[(C\Omega + D) \frac{\partial}{\partial Z} \left\{ (C\bar{\Omega} + D) \frac{\partial f}{\partial \bar{Z}} \right\} \right] \\ &= \det(Y) |\det(C\Omega + D)|^{-2} \det \left[(C\Omega + D) \left\{ (C\bar{\Omega} + D) \left(\frac{\partial}{\partial Z} {}^t \left(\frac{\partial f}{\partial \bar{Z}} \right) \right) \right\} \right] \\ &= \det(Y) |\det(C\Omega + D)|^{-2} \det \left[(C\Omega + D) \frac{\partial}{\partial Z} \left(\frac{\partial f}{\partial \bar{Z}} \right) {}^t (C\bar{\Omega} + D) \right] \\ &= \det(Y) \det \left(\frac{\partial}{\partial Z} {}^t \left(\frac{\partial f}{\partial \bar{Z}} \right) \right) \\ &= \mathbb{K} f, \end{aligned}$$

where $\mathbb{K}_*$ denotes the image of $\mathbb{K}$ under the transformation

$$(\Omega, Z) \longmapsto (\Omega_*, Z_*) = ((M \cdot \Omega, (Z + \lambda\Omega + \mu)(C\Omega + D)^{-1})$$

with $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in Sp(n, \mathbb{R})$ and $(\lambda, \mu; \kappa) \in H_{\mathbb{R}}^{(n,m)}$.

From (3.35), (3.40) and the fact that the action (1.2) of G^J on $\mathbb{H}_{n,m}$ is compatible with the action (3.31) of G_*^J on $\mathbb{D}_{n,m}$ via the partial Cayley transform, we see that

$$(3.43) \quad \mathbb{K}_{\mathbb{D}_{n,m}} = \det(I_n - W\bar{W}) \det \left(\frac{\partial}{\partial \eta} {}^t \left(\frac{\partial}{\partial \bar{\eta}} \right) \right)$$

is a G_*^J -invariant differential operator on $\mathbb{D}_{n,m}$.

Using (3.21), (3.22) and Lemma 3.1, we can show without difficulty that the matrix-valued differential operator $\mathbb{T}$ on $\mathbb{H}_{n,m}$ defined by

$$(3.44) \quad \mathbb{T} = \left(\frac{\partial}{\partial \bar{Z}} \right) {}^t Y \frac{\partial}{\partial Z}$$

is invariant under the action (1.2) of G^J . Therefore each (k, l) -entry $\mathbb{T}_{kl}$ of $\mathbb{T}$ given by

$$(3.45) \quad \mathbb{T}_{kl} = \sum_{i,j=1}^n y_{ij} \frac{\partial^2}{\partial \bar{z}_{ki} \partial z_{lj}}, \quad 1 \leq k, l \leq m$$

is an element of $\mathbb{D}(\mathbb{H}_{n,m})$. From (3.35), (3.40) and the fact that the action (1.2) of G^J on $\mathbb{H}_{n,m}$ is compatible with the action (3.31) of G_*^J on $\mathbb{D}_{n,m}$ via the partial Cayley

transform, we see that

$$(3.46) \quad \mathbb{T}^{\mathbb{D}} := {}^t \left(\frac{\partial}{\partial \bar{\eta}} \right) (I_n - \bar{W}W) \frac{\partial}{\partial \eta}$$

is a G_*^J -invariant matrix-valued differential operator on $\mathbb{D}_{n,m}$. Thus each (k, l) -entry $\mathbb{T}_{kl}^{\mathbb{D}}$ of $\mathbb{T}^{\mathbb{D}}$ given by

$$(3.47) \quad \mathbb{T}_{kl}^{\mathbb{D}} = \sum_{i,j=1}^n \left(\delta_{ij} - \sum_{r=1}^n \bar{w}_{ir} w_{jr} \right) \frac{\partial^2}{\partial \bar{\eta}_{ki} \partial \eta_{lj}}, \quad 1 \leq k, l \leq m$$

is a G_*^J -invariant differential operator on $\mathbb{D}_{n,m}$.

Indeed it is very complicated and difficult at this moment to express the generators of the algebra of all G_*^J -invariant differential operators on $\mathbb{D}_{n,m}$ explicitly. In particular, it is extremely difficult to find explicit G_*^J -invariant differential operators on $\mathbb{D}_{n,m}$ of *odd* degree. We propose an open problem to find other explicit G_*^J -invariant differential operators on $\mathbb{D}_{n,m}$.

Example 3.2. We consider the case $n = m = 1$. For a coordinate (ω, z) in $T_{1,1}$, we write $\omega = x + i y$, $z = u + i v$, x, y, u, v real. Then the algebra $\text{Pol}_{1,1}^{U(1)}$ is generated by

$$\begin{aligned} q(\omega, z) &= \frac{1}{4} \omega \bar{\omega} = \frac{1}{4} (x^2 + y^2), \\ \xi(\omega, z) &= z \bar{z} = u^2 + v^2, \\ \phi(\omega, z) &= \frac{1}{2} \text{Re}(z^2 \bar{\omega}) = \frac{1}{2} (u^2 - v^2)x + uv y, \\ \psi(\omega, z) &= \frac{1}{2} \text{Im}(z^2 \bar{\omega}) = \frac{1}{2} (v^2 - u^2)y + uv x. \end{aligned}$$

We put

$$D_1 = \Theta(q), \quad D_2 = \Theta(\xi), \quad D_3 = \Theta(\phi) \quad \text{and} \quad D_4 = \Theta(\psi).$$

Using Formula (3.3), we can show that the algebra $\mathbb{D}(\mathbb{H}_{1,1})$ is generated by the following differential operators

$$\begin{aligned} D_1 &= y^2 \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) + v^2 \left(\frac{\partial^2}{\partial u^2} + \frac{\partial^2}{\partial v^2} \right) \\ &\quad + 2 y v \left(\frac{\partial^2}{\partial x \partial u} + \frac{\partial^2}{\partial y \partial v} \right), \\ D_2 &= y \left(\frac{\partial^2}{\partial u^2} + \frac{\partial^2}{\partial v^2} \right), \end{aligned}$$

$$D_3 = y^2 \frac{\partial}{\partial y} \left(\frac{\partial^2}{\partial u^2} - \frac{\partial^2}{\partial v^2} \right) - 2y^2 \frac{\partial^3}{\partial x \partial u \partial v} \\ - \left(v \frac{\partial}{\partial v} + 1 \right) D_2$$

and

$$D_4 = y^2 \frac{\partial}{\partial x} \left(\frac{\partial^2}{\partial v^2} - \frac{\partial^2}{\partial u^2} \right) - 2y^2 \frac{\partial^3}{\partial y \partial u \partial v} \\ - v \frac{\partial}{\partial u} D_2,$$

where $\tau = x + iy$ and $z = u + iv$ with real variables x, y, u, v . Moreover, we have

$$D_1 D_2 - D_2 D_1 = 2y^2 \frac{\partial}{\partial y} \left(\frac{\partial^2}{\partial u^2} - \frac{\partial^2}{\partial v^2} \right) \\ - 4y^2 \frac{\partial^3}{\partial x \partial u \partial v} - 2 \left(v \frac{\partial}{\partial v} D_2 + D_2 \right).$$

In particular, the algebra $\mathbb{D}(\mathbb{H}_{1,1})$ is not commutative. We refer to [1, 21] for more detail.

REFERENCES

- [1] R. Berndt and R. Schmidt, *Elements of the Representation Theory of the Jacobi Group*, Progress in Mathematics, **163**, Birkhäuser, Basel, 1998.
- [2] D. Bump and Y. J. Choie, *Derivatives of modular forms of negative weight*, Pure Appl. Math. Q. **2** (2006), no. 1, 111-133.
- [3] M. Eichler and D. Zagier, *The Theory of Jacobi Forms*, Progress in Mathematics, **55**, Birkhäuser, Boston, Basel and Stuttgart, 1985.
- [4] Harish-Chandra, *Representations of a semisimple Lie group on a Banach space. I.*, Trans. Amer. Math. Soc. **75** (1953), 185-243.
- [5] Harish-Chandra, *The characters of semisimple Lie groups*, Trans. Amer. Math. Soc. **83** (1956), 98-163.
- [6] S. Helgason, *Differential operators on homogeneous spaces*, Acta Math. **102** (1959), 239-299.
- [7] S. Helgason, *Groups and geometric analysis*, Academic Press, New York (1984).
- [8] R. Howe, *Perspectives on invariant theory: Schur duality, multiplicity-free actions and beyond*, The Schur lectures (1992) (Tel Aviv), Israel Math. Conf. Proceedings, **vol. 8** (1995), 1-182.
- [9] H. Maass, *Die Differentialgleichungen in der Theorie der Siegelschen Modulfunktionen*, Math. Ann. **126** (1953), 44-68.
- [10] H. Maass, *Siegel modular forms and Dirichlet series*, Lecture Notes in Math., **vol. 216**, Springer-Verlag, Berlin-Heidelberg-New York (1971).
- [11] I. Satake, *Algebraic Structures of Symmetric Domains*, Kano Memorial Lectures 4, Iwanami Shoten, Publishers and Princeton University Press (1980).
- [12] G. Shimura, *Invariant differential operators on hermitian symmetric spaces*, Ann. Math. **132** (1990), 237-272.

- [13] C. L. Siegel, *Symplectic Geometry*, Amer. J. Math. **65** (1943), 1-86; Academic Press, New York and London (1964); Gesammelte Abhandlungen, no. 41, vol. II, Springer-Verlag (1966), 274-359.
- [14] C. L. Siegel, *Topics in Complex Function Theory: Abelian Functions and Modular Functions of Several Variables*, vol. III, Wiley-Interscience, 1973.
- [15] H. Weyl, *The classical groups: Their invariants and representations*, Princeton Univ. Press, Princeton, New Jersey, second edition (1946).
- [16] J.-H. Yang, *The Siegel-Jacobi Operator*, Abh. Math. Sem. Univ. Hamburg **63** (1993), 135-146.
- [17] J.-H. Yang, *Singular Jacobi Forms*, Trans. Amer. Math. Soc. **347** (6) (1995), 2041-2049.
- [18] J.-H. Yang, *Construction of vector valued modular forms from Jacobi forms*, Canad. J. Math. **47** (6) (1995), 1329-1339 or arXiv:math.NT/0612502.
- [19] J.-H. Yang, *A geometrical theory of Jacobi forms of higher degree*, Proceedings of Symposium on Hodge Theory and Algebraic Geometry (edited by Tadao Oda), Sendai, Japan (1996), 125-147 or Kyungpook Math. J. **40** (2) (2000), 209-237 or arXiv:math.NT/0602267.
- [20] J.-H. Yang, *The Method of Orbits for Real Lie Groups*, Kyungpook Math. J. **42** (2) (2002), 199-272 or arXiv:math.RT/0602056.
- [21] J.-H. Yang, *A note on Maass-Jacobi forms*, Kyungpook Math. J. **43** (4) (2003), 547-566 or arXiv:math.NT/0612387.
- [22] J.-H. Yang, *Invariant metrics and Laplacians on Siegel-Jacobi space*, Journal of Number Theory **127** (2007), 83-102 or arXiv:math.NT/0507215.
- [23] J.-H. Yang, *A partial Cayley transform for Siegel-Jacobi disk*, J. Korean Math. Soc. **45**, No. 3 (2008), 781-794 or arXiv:math.NT/0507216.
- [24] J.-H. Yang, *A note on a fundamental domain for Siegel-Jacobi space*, Houston J. Math. **32** (3) (2006), 701-712 or arXiv:math.NT/0507218.
- [25] J.-H. Yang, *Invariant metrics and Laplacians on Siegel-Jacobi disk*, arXiv:math.NT/0507217 or Chinese Annals of Mathematics, DOI:10.1007/s11401-008-0348-7, Springer-Verlag Berlin Heidelberg (2009).

DEPARTMENT OF MATHEMATICS, INHA UNIVERSITY, INCHEON 402-751, KOREA

E-mail address: jhyang@inha.ac.kr